


XV. *Of the Penetration of a Hemisphere by an indefinite Number of equal and similar Cylinders.* By Thomas Knight, Esq.
Communicated by Sir Humphry Davy, LL. D. Sec. R. S.

Read March 19, 1812.



ing certain portions of the surface and solidity of a hemisphere, form, together, a single case of the following problem; which is one of the most remarkable, for generality and simplicity of result, in the whole compass of geometry.

Problem.

To pierce a hemisphere, perpendicularly on the plane of its base, with any number of equal and similar cylinders; of such a kind, that, if we take away from the hemisphere those portions of the cylinders that are within it, the remaining part shall admit of an exact cubature: and if we take away, from the surface of the hemisphere, those portions cut out by the cylinders, the remaining surface shall admit of an exact quadrature.

Let fig. 1 represent the nearest half of the hemisphere, where a is the pole, bdcf a quadrant of the great circle form-

ing its base. From every point d , on this side of b , draw the radius dC to the centre of the hemisphere, and (if the number of cylinders is to be $2n^*$) take the arc bs equal to n times the arc bd , draw se perpendicular to Cb , and with the centre C and radius Ce describe the arc er cutting Cd in r . Through all the points (r) thus found, draw the curve line brC , terminated at b and C , and it shall be half the base of one of the required cylinders.

It is, in the first place, evident, from the construction, that the half cylinder, whose base is $beCrb$, is contained between two planes $CabC$, $CacC$, making with each other the angle $bCc = \frac{90^\circ}{n}$; consequently the whole base of the hemisphere may be pierced by $2n$ such cylinders as this is the half of.

Let $atmb$ be the intersection of the surfaces of the half cylinder and hemisphere; and a great circle passing through a and d , and meeting $atmb$ at m . Call the radius of the sphere r , Cr is the cosine of the arc bs to the radius r , by construction; it is also the cosine of the arc md to the same radius; therefore $md = bs = n \times bd$.

Put $bd = \phi$; $md = n \times \phi$; $d\phi = \psi$. Moreover, put A for the spherical space $atmbdcna$ contained by the arcs anc , cdb and the curve $atmb$; and let S be the solidity of the portion of the hemisphere contained between the quadrant $ancC$ and the surface ($brCatmb$) of the half cylinder. It is easy to see that

$$\begin{aligned} A &= r^3 \iint \phi \cos. \psi \times \psi, \\ S &= \frac{r^3}{2} \iint \phi \cos. \psi \times \cos. \psi \times \cos. \psi \psi \\ &\quad - \frac{r^3}{2} \iint \phi \cos. n\phi \times \cos. n\phi \times \cos. \psi \psi. \end{aligned}$$

* I do not intend $2n$ to represent an even number *only*, n may be $\frac{1}{2}$, or $\frac{3}{2}$, or $\frac{5}{2}$, &c. and $2n$ express any number whatever.

The fluents to be taken, first from $\psi = 0$ to $\psi = n\phi$, and then from $\phi = 0$, to $\phi = \frac{90^\circ}{n}$. The first operation gives

$$A = r^2 \int \phi \sin. n\phi,$$

$$S = \frac{r^3}{2} \int \phi \left\{ \frac{3}{4} \sin. n\phi + \frac{1}{12} \sin. 3n\phi \right\} - \frac{r^3}{2} \int \phi \sin. n\phi \cos. 2n\phi,$$

and by the second we get

$$A = C - \frac{r^2}{n} \cos. n\phi$$


$$S = \frac{r^3}{2n} \left\{ \frac{\cos. 3n\phi}{3} - \frac{3}{4} \cos. n\phi - \frac{1}{36} \cos. 3n\phi \right\} + C,$$

which fluents being taken from $n\phi = 0$, to $n\phi = 90^\circ$, are

$A = \frac{r^2}{n}$; $S = \frac{2}{9} \times \frac{r^3}{n}$; and if these are multiplied by $4n$, we have

$$A = 4r^2; S = \frac{8}{9} r^3;$$

for the whole that remains of the surface and solidity of the hemisphere after the subduction of the $2n$ cylinders. Thus A and S (for the whole hemisphere) do not depend on the number of the cylinders with which the penetration is made; *a most remarkable circumstance, seeing that amongst the bases of those cylinders are curves of an infinity of different kinds and orders.*

Let fig. 2 represent half the base of one of the cylinders;

 Cb the radius of the hemisphere, C the centre. From r , any point in the curve, let fall the perpendicular rp on the axis; call Cp , x ; rp , y .

By construction, $Cr = \sqrt{x^2 + y^2} = r \cos. n \cdot bCr$; now the cosine of the simple arc bCr is $\frac{x}{\sqrt{x^2 + y^2}}$, which being put in the trigonometrical expression for the cosine of the multiple arc

in terms of the cosine of the simple one, we have, for the equation of the curve brC.

When $n = 1$, $\sqrt{x^2 + y^2} = \frac{rx}{\sqrt{x^2 + y^2}}$ or $x^2 + y^2 = rx$, the equation of a circle.

When $n = 2$, $\sqrt{x^2 + y^2} = \frac{2rx^2}{x^2 + y^2} - r$ or $(x^2 + y^2)^3 = r^2(x^2 - y^2)^2$; and in general the curve will be algebraic when n is any whole number.