

XVII. *On the Meteorology of the Lake District of Cumberland and Westmoreland ; including the results of Experiments on the fall of Rain at various heights, up to 3166 feet above the sea level.*

By JOHN FLETCHER MILLER, *Esq.*, *F.R.A.S.*

Communicated by Lieut.-Col. SABINE, *For. Sec. R.S.*

Received February 19,—Read April 19, 1849.

THE form of the instruments, their positions, &c., are fully described in a former paper, read before the Royal Society on the 18th of May 1848.

The Roman numerals attached to each gauge refer to corresponding numbers on a map of the Lake District accompanying the said paper.

With one exception, elsewhere referred to, no accident of any importance has occurred to interrupt the continuity or correctness of the rain tables. Indeed, considering the extent of the experiments, and the exposed position of many of the instruments, mishaps of any kind have from the first been exceeding rare. The gauges are all under my own supervision ; the registrars are all thoroughly instructed in the method of reading off the rain, which is regularly recorded in registers prepared for the purpose ; and, being all careful persons, and almost constantly on the spot, the observations are seldom omitted even for a single day.

The records are transmitted to me at the close of each month, and are entered in collateral columns in a large folio journal ; should there be any apparent discrepancy at any station on any particular day or days, an explanation is requested, and the original register examined, to ascertain whether the transmitted copy is correct. In this way, errors have occasionally, though very rarely, been discovered, and I do not recollect that one such has occurred during the past year.

The Observatory, Whitehaven,
February 6, 1849.

TABLE I.—Synopsis of the Fall of Rain in the Lake Districts of Cumberland and Westmoreland, for the year 1848.

No.	I.	II.	III.	IV.	V.	VI.	VII.	VIII.	IX.	X.	XI.	XII.	XIII.	XIV.	XV.	XVI.	XVII.	XVIII.	XIX.	XX.	Prevailing winds.			
1848.	Whitehaven.			The Flish, three miles south of Whitehaven.	Cockermouth.	Passenthwaite Halls, 210 feet above the sea.	Kewick, 258 feet above the sea.	Vale of Gillerthwaite, Innerdale, 280 feet above the sea.	Loweswater, 336 feet above the sea.	Foot of Crummock Lake, 283 feet above the sea.	Gatesgarth, 326 feet above the sea.	Eskdale.		Wastdale Head, 166 feet above the sea.	Westmoreland.			Borrowdale.		Stonehwaite.	At Crummock Lake, two observations daily.	At Seathwaite, one observation daily.	At the coast, two observations daily.	
	High Street, 90 feet above the sea.	Round Close, 180 feet above the sea.	St. James's Church Steeple, 78 feet above the sea.									Centre of Vale.	Head of Vale.		The How, Troutbeck, 300 feet above the sea.	Ambleside, 190 feet above the sea*.	Langdale Head, 250 feet above the sea.	In Garden, 6 inches above the surface.	In Field, 18 inches above the surface.					
January	in. 3745	in. 3735	in. 3503	in. 434	in. 380	in. 342	in. 4112	in. 720	in. 5184	in. 730	in. 825	in. 492	in. 635	in. 606	in. 5922	in. 472	in. 701	in. 967	in. 963	in. 786	E.	variable.	S.E. & S.W.	
February	7815	7986	6397	982	973	811	10276	1693	12971	1448	2478	1170	1422	1992	19903	1419	2567	3055	2998	2290	886	S.W.	W.	S.W. & W.
March	4588	3930	3498	538	380	300	4736	730	5706	677	857	556	670	915	6834	585	971	1136	1118	886	S.W.	W. & S.W.	N.E.	
April	495	339	293	87	100	223	1722	300	1627	283	354	187	282	324	1927	230	332	419	412	349	E.	variable.	N.E.	
May	1798	1612	1452	223	71	53	1062	340	1175	152	317	101	183	154	1852	203	211	305	297	303	S.W.	W.	S.W.	
June	3867	4041	3043	524	436	353	5036	616	6236	717	831	643	634	987	6478	574	986	1130	1119	982	S.W.	S.W.	S.W.	
July	3630	3862	2442	460	410	345	6246	958	7431	1041	1366	598	721	1066	6080	618	1212	1776	1753	1434	S.W.	S.W.	W.	
August	5054	5501	4188	807	731	472	6280	926	7795	973	1137	745	842	1074	7488	716	1141	1391	1354	1036	S.W.	S.W.	W.	
September ..	2266	2122	1723	272	251	221	2104	486	3755	463	543	372	399	547	3257	280	484	700	692	488	S.W. var.	S.W. var.	S.W.	
October	5772	5676	4632	836	444	512	7158	970	8134	1111	1572	787	990	1188	9357	989	1353	1732	1681	1427	S.	S. var.	S.W.	
November...	3507	3412	2186	407	480	477	6703	960	6552	762	1240	613	797	1110	9954	666	1313	1407	1354	1138	N.W. var.	N.W. & S.W.	N.W. var.	
December...	4805	4484	3187	512	581	597	10972	1074	10102	1450	1835	774	1103	1569	12295	1006	1767	2071	1981	1905	S. & S.E.	S.W.	S.W.	
1848.	47342	46700	36344	6082	5237	4706	66407	9773	76688	9807	13355	7038	8678	11532	91347	7758	13038	16089	13722	13024	S.W.	S.W.	S.W.	
1847.	42921	42023	30713	4780	4255	4445	58286	8013	66296	8232	10625	5866	7493	9634	78004	11295	12924	12680	10621	westerly	S.W.	S.W.	
1846.	49134	35422	5516	5241	67678	8387	79249	9647	12190	10693	77719	12740	14351	
1845.	49207	33489	5300	4693	62202	7688	69542	8748	12413	10855	76305	13600	15187	
1844.	
1843.	18561	14550	1987	21781	2746	25650	3071	3944	4151	25576	
July to December																								

* The gauge at Grasmere, which in the Tables and Map accompanying the Report for 1847 was marked No. XVI., has been discontinued, and Ambleside is inserted in its place.

TABLE II.
Wet Days.

1848.	Whitehaven.	The Fosh.	Cockermouth.	Bassenthwaite Halls.	Keswick.	Loweswater Lake.	Crummock Lake.	Eskdale.	Wastdale Head.	Troutbeck.	Langdale Head.	Seathwaite.	Stonethwaite.
January ...	13	13	15	14	15	10	12	11	15	7	13	14	14
February...	23	23	25	25	23	25	23	22	24	23	23	25	23
March.....	23	22	25	19	26	26	23	24	26	26	24	24	24
April	16	12	16	13	16	13	12	14	18	13	13	16	16
May	6	7	6	4	10	8	6	6	8	10	7	11	11
June	19	19	20	17	20	18	19	19	22	18	20	19	21
July.....	18	18	19	15	17	18	17	17	21	16	18	19	18
August ...	19	23	23	18	23	23	24	23	25	24	24	25	22
September.	13	11	15	11	10	14	14	10	16	10	12	16	13
October ...	22	22	24	26	27	23	21	21	24	18	21	22	23
November..	20	20	22	18	24	21	18	21	23	18	19	22	21
December..	18	17	18	16	18	18	18	17	21	18	18	19	18
1848.	210	207	228	196	229	217	207	205	243	201	212	232	224
1847.	191	183	210	199	204	190	199	226	188	209	202	195
1846.	200	208	234	213	198	216	234	194	213	219	
1845.	193	175	212	195	195	202	211	180	211	

TABLE III.

Showing the Quantity of Rain received by the Mountain Gauges in thirteen months, between the 1st of December 1847, and the 31st of December 1848.

No.	XXI.	XXII.	XXIII.	XXIV.	XXV.	XIV.	XIII.	XXVI.	XIX.
1848.	Sea Fell, 3166 feet above the sea.	Great Gabel, 2925 feet above the sea.	Sparkling Tarn, 1900 feet above the sea.	Stye Head, 1290 feet above the sea.	Brant Rigg, about 500 feet above the sea.	Valley.		Borrowdale.	
						To the west, Wastdale, 166 feet above the sea.	To the south-east, Eskdale, height unknown.	Seatollar Common, 1334 feet above the sea.	The Valley, Seathwaite, 242 feet above the sea.
	in.	in.	in.	in.	in.	in.	in.	in.	in.
December..	Frozen	Frozen	Frozen.	Frozen	12·15	8·93	12·53	20·33
1848.	Frozen	Frozen	Frozen.	Frozen	6·06	6·35	Frozen.	9·63
January	Frozen	39·45	41·31	32·65	19·92	14·22	34·18	29·98
February...	28·00	8·04	9·17	9·36	9·15	6·70	7·10	11·18
March.....	2·29	2·31	3·22	5·84	3·24	2·82	2·07	4·12
April	2·60	3·72	3·10	1·98	1·54	1·83	2·80	2·97
May	*4·42	9·00	11·52	9·40	7·80	9·87	6·34	10·00	11·19
June	8·58	10·32	17·40	13·18	8·12	10·66	7·21	15·00	17·53
July.....	10·65	9·79	13·09	12·31	9·16	10·74	8·42	11·55	13·54
August ...	9·74	5·39	8·16	7·51	4·76	5·47	3·99	5·42	6·92
September..	5·11	9·71	17·06	14·85	11·36	11·88	9·90	13·20	16·81
October ...	10·96	6·17	11·44	11·34	9·36	11·10	7·97	12·63	13·54
November..	6·81	8·05	16·40	13·33	8·80	15·69	11·03	13·00	19·81
December..	8·46								
Inches... {	64·73 From May }	91·32	148·59	138·72	109·19	127·47	95·71	139·48	177·55

* The fall of rain at this station, during the winter of 1847-48, was lost, in consequence of injury sustained by the receiver from the frost.

TABLE IV.—For the Summer Months.

No.	XXI.	XXII.	XXIII.	XXIV.	XXV.	XIV.	XIII.	XXVI.	XIX.
1848.	Sca Fell, 3166 feet above the sea.	Great Gabel, 2925 feet above the sea.	Sparkling Tarn, 1900 feet above the sea.	Stye Head, 1290 feet above the sea.	Brant Rigg, about 500 feet above the sea.	Valley.		Borrowdale.	
						To the west of Wast- dale	To the south-east of Eskdale.	Seatollar Common, 1334 feet above the sea.	The Valley, Seathwaite, 242 feet above the sea.
May	in. 4·42	in. 2·60	in. 3·72	in. 3·10	in. 1·98	in. 1·54	in. 1·83	in. 2·80	in. 2·97
June	8·58	9·00	11·52	9·40	7·80	9·87	6·34	10·00	11·19
July	10·65	10·32	17·40	13·18	8·12	10·66	7·21	15·00	17·53
August	9·74	9·79	13·09	12·31	9·16	10·74	8·42	11·55	13·54
September	5·11	5·39	8·16	7·51	4·76	5·47	3·99	5·42	6·92
October	10·96	9·71	17·06	14·85	11·36	11·88	9·90	13·20	16·81
Inches.....	49·46	46·81	70·95	60·35	43·18	50·16	37·69	57·97	68·96

TABLE V.—For the Winter Months.

1847.	Sea Fell.	Great Gabel.	Sparkling Tarn.	Stye Head.	Brant Rigg.	The Valley.		Borrowdale.	
						To the west of Wast- dale.	To the south-east of Eskdale.	Seatollar Common.	Valley. Seathwaite.
December	in.	in. Frozen	in. Frozen	in. Frozen	in. Frozen	in.	in.	in. 12·53	in. 20·33
1848. January	Frozen	Frozen	Frozen	Frozen	Frozen	9·63
February	Frozen	39·45	41·31	32·65	38·13	29·50	34·18	29·98
March.....	28·00	8·04	9·17	9·36	9·15	6·70	7·10	11·18
April	2·29	2·31	3·22	5·84	3·24	2·82	2·07	4·12
November	6·81	6·17	11·44	11·34	9·36	11·10	7·97	12·63	13·54
December	8·46	8·05	16·40	13·33	8·80	15·69	11·03	13·00	19·81
Inches.....	44·51	77·64	78·37	66·01	77·31	58·02	81·51	108·59

TABLE VI.—Temperature at Seathwaite taken by Self-registering Thermometers made by WATKINS and HILL.

1848.	Absolute.		Mean of max.	Mean of min.	Approximate mean tempera- ture.	Mean at 9 A.M.	On grass.				Prevailing winds.
	Max.	Min.					Min.	Mean.	Radiation.		
									Max.	Mean.	
January	50	9	37.43	27.50	32.46	32.71	6.5	24.56	7	2.93	Var.
February	49.5	25	44.50	36.67	40.58	40.24	19	33.74	7	2.93	w.
March.....	57.5	27	46.66	34.14	40.40	39.67	19.5	29.44	8.5	4.81	w. and s.w.
April	67	28.5	52.78	38.03	45.40	45.35	21	32.38	10.5	5.65	Var.
May	72.5	32	63.52	48.20	55.86	55.89	24	38.40	15	9.80	w.
June	75	40	63.86	50.75	57.30	57.18	29	42.36	15	8.41	s.w.
July	79	45	62.61	53.32	57.96	57.19	32	46.27	14	7.05	s.w.
August	62.5	41.5	57.99	49.87	53.93	53.48	29	40.79	15.5	9.08	s.w.
September	68	38.5	57.93	49.05	53.49	52.28	25.5	39.76	16	9.28	s.w. var.
October	60.5	29.5	50.17	43.66	46.91	46.61	18	36.01	14.5	7.96	s. var.
November	52	24	44.59	36.35	40.47	40.13	12	29.63	13.5	6.72	n.w. and s.w.
December	56	26	43.76	37.21	40.48	40.43	11	28.86	18.7	8.38	s.w.
Means at Whitehaven }	62.4 62.5	30.5 33.0	52.15 53.62	42.06 44.00	47.10 48.81	46.76	20.5 23.1	35.18 37.24	12.9	6.91 6.76	s.w. Var.
Difference	0.1	2.5	1.47	1.94	1.71	2.6	2.06	0.15	
Difference in 1847..... }	0.5	3.8	0.96	1.46	1.22						

Remarks.

The fall of rain in the Lake District during the year 1848, greatly exceeds the amount in any other year since the register was commenced in 1844; and the same remark applies to the number of wet days. The total depth in 1848, at Seathwaite, the wettest station, is 160.89 inches, and of this quantity, 114.32 inches fell in the six months comprehending February, July, August, October, November and December.

The wettest quarter of the year was the last, in which 52.10 inches were measured; the wettest month in 1848 was February, which yielded 30.55 inches, by far the largest quantity ever measured in any month in this country; and the two wettest days were the 3rd and 26th of December, when 4.60 and 4.22 inches respectively, were read off.

At Seathwaite, there have been forty-eight days in last year wherein the quantity of rain fallen was between half an inch and 1 inch; thirty-two days between 1 and 2 inches; thirteen days between 2 and 3 inches; five days between 3 and 4 inches; and two days between 4 and 5 inches.

In a former paper which I had the honour to lay before the Royal Society, I endeavoured to give a general outline of the meteorology of the Lake District, as far as the facts then ascertained would permit. I now proceed to discuss one or two points which were intentionally passed over in that report.

The mountains flanking the Lake District valleys, generally increase in altitude

with great regularity, towards the head or eastern extremity of the vale; and it is here that the greatest depth of rain is invariably found. The difference in the annual quantity between places contiguous to each other and in the same valley, is often remarkably great. The amount increases rapidly as we recede from the sea, and towards the head of the valley the incremental ratio is enormous.

Loweswater, Buttermere and Gatesgarth, in the same line of valley, are about two miles apart from each other; yet in 1848 Loweswater has received 76 inches, Buttermere 98 inches, and Gatesgarth $133\frac{1}{2}$ inches of water. Here in a space of four miles, we have a difference of 57 inches in twelve months, and in some years the proportional excess is still greater. The head of Eskdale receives fully one-fourth more rain than the middle of the valley, and a like difference obtains between two stations in the Vale of Borrowdale about a mile apart, whilst the proportion between the deposit at Ennerdale Lake and a farm-house three miles to the westward, is as two to one nearly.

At an early stage of this inquiry, I was forcibly struck with the rapid increment in the fall towards the head or terminal point of all valleys, and I made some experiments in order to ascertain whether the effect was appreciable at much shorter distances than any of those just referred to. For this purpose I caused a duplicate gauge to be made, in all respects exactly similar to the one at Wastdale Head, and fixed it about 200 yards higher up the valley. The two gauges were read off daily at the same hour for twelve months, and the following are the results:—

Wastdale Head.

1845.	No. 1.	No. 2.	Difference.
	in.	in.	in.
October	12·35	11·89	0·46
November	12·31	11·90	0·41
December	16·18	15·78	0·40
1846.			
January	12·97	12·47	0·50
February	6·60	6·58	0·02
March	10·35	10·07	0·28
April	6·59	6·16	0·43
May	3·65	3·44	0·21
June	5·33	4·88	0·45
July	16·82	16·59	0·23
August	8·96	8·97	—0·01
September	3·79	3·64	0·15
Inches	115·90	112·37	3·53

It will be observed that the higher gauge, marked No. 1, is always in excess, and that the difference in a single month sometimes amounts to half an inch, though the instruments are within two or three fields' breadth of each other. Here the effect of a slight increase in proximity to the higher mountains is very apparent.

Temperature.—The mountain valleys are commonly supposed to be intensely cold, particularly in the winter season; but the thermometer, so far from countenancing

this opinion, shows that the inhabitants enjoy a milder and more equable climate than those who reside in the open country. The town of Whitehaven, from its proximity to the sea on the west coast, is well known to have a much higher mean temperature than is due to its latitude; it is also much less subject to those great and sudden fluctuations of heat and cold to which inland places are liable. Yet the mean temperature at Seathwaite, in the heart of the lake country, is only about $1^{\circ}5$ lower than with us. In 1847 and 1848 the mean temperature of Seathwaite was $47^{\circ}46$ and $47^{\circ}10$, whilst at Kendal it was $46^{\circ}67$ and $46^{\circ}32$ respectively.

In winter, the mean of the night temperature is several degrees higher than at Cockermouth in the open plain, where the frost is much more severe. The indications of the thermometer are in accordance both with the assertions of the residents and with my own observation; for in travelling to the lakes, where the roads over the commons were frozen hard, I have often found them quite soft and clammy on arriving amongst the hills.

These valleys not only have a higher winter temperature than many localities greatly to the south of them, but they very rarely experience those low extremes which not unfrequently occur in the southern counties of England. The mean temperature of the winter months at Chiswick, in Middlesex, is nearly the same as in the Lake District, whilst a much greater extreme of cold is frequently felt there than in the north. In the neighbourhood of the metropolis the thermometer sometimes indicates a degree of cold almost unknown in these districts. Thus, on the night between the 11th and 12th of February 1847, the temperature at Greenwich fell to 6° , at Chiswick to 4° , and at Uckfield, Sussex, to 1° ; when at Seathwaite the minimum was $24^{\circ}5$, and the minimum for the month 20° .

The lakes, by absorbing heat in the summer and giving it out in the winter months, added to that radiated from the rocky mountain breasts, and, above all, the caloric evolved in a sensible form by the condensation of such enormous volumes of vapour, no doubt tend greatly to modify the climate of these sequestered localities.

Temperature on Sca Fell.—Last summer I stationed a pair of RUTHERFORD'S self-registering thermometers (previously compared with a standard) on the top of Sca Fell Pike; they are suspended in a deal box, having the sides and base riddled with small circular holes, so that the instruments are freely exposed to the air, and at the same time thoroughly protected from the effects of terrestrial radiation. On the summit of the Pike is a cairn, or large pile of stones about 8 feet in height, having a stout pole in the centre, which projects about 2 feet above the top of the pile. To this pole the box containing the thermometers is firmly fixed.

From the maximum thermometer I have never been able to obtain any correct readings, as, from some cause, the steel needle is always found at the extreme end of the stem, furthest from the bulb. I cannot account for this, unless indeed the fine steel needle is affected by electrical currents at such an extreme height in the clouds. The readings of the maximum thermometer would, however, have probably been of

little value, as it would be almost impossible to protect it from the effect of solar radiation.

The following are the readings of the minimum thermometer for each month from July to the end of the year 1848 :—

July, 22°; August, 24°; September, 18°; October, —6°; November, —6°; December, —9°, or 41° below the freezing-point of water*.

The lowest extreme in these months, in the Vale of Borrowdale, at 4 feet above the ground, was as under :—

July, 45°; August, 41°·5; September, 38°·5; October, 29°·5; November, 24°; December, 26°.

The Mountain Gauges.—The results are in strict accordance with those of the two previous years, and confirm the correctness of the conclusion drawn from them in a former paper, “that the quantity of rain increases from the valley upwards to an altitude of about 2000 feet, above which it begins to diminish.”

Thus, in thirteen months,—

	Inches.
The Valley . . . 160 feet above sea, has received	127·47
Stye Head . . . 1290 feet above sea, has received	138·72
Seatollar Common 1334 feet above sea, has received	139·48
Sparkling Tarn . 1900 feet above sea, has received	148·59
Great Gabel . . 2925 feet above sea, has received	91·32

I regret to state that the whole quantity of water collected in the Sca Fell gauge during the winter of 1847–48 was lost, in consequence of injury caused by the frost. In the spring of last year I had a new set of receivers constructed for these stations, which are made of very heavy sheet copper, double-lapped at the seams, and with the bottoms convex inwards, to enable them the better to bear the expansive force of the water during its conversion into ice; so that a similar accident is not likely to occur again.

From the table for the summer months, it appears that between the 1st of May and the 31st of October, the gauge at 1290 feet has received $20\frac{1}{2}$ per cent. more rain than the valley; at 1334 feet, $15\frac{1}{2}$ per cent. more; at 1900 feet, $41\frac{1}{2}$ per cent. more; at 2928 feet, 6 per cent. *less*; and at 3166 feet, about 1 per cent. *less* than the valley. The excess over the valley is somewhat greater at all the stations than in the two previous years, and Sca Fell, which usually obtains *less* rain than Gabel, has this summer received *more*.

By referring to the table for the winter months, we find that the station at 1290 feet has obtained 0·5 per cent. more rain than the valley; at 1334 feet, $5\frac{1}{2}$ per cent.

* On the 29th and 31st of January 1849, the box containing the thermometers was so thickly encased in ice, that it could not be opened. The minimum temperature for the month was read off on the 12th of February, being no less than 34° below the zero point of FAHRENHEIT'S scale. This unheard-of extreme of cold undoubtedly occurred on the night between the 2nd and 3rd of January, when a naked thermometer on grass, at Whitehaven, fell to +4°, and one on raw wool to —2°·8.—J. F. M.

more; at 1900 feet, $1\frac{1}{2}$ per cent. more; and at 2928 feet, $42\frac{1}{2}$ per cent. *less* than the valley. Here the gauge at 1334 feet, which on the average of the two preceding winters received the same quantity as the Vale of Wastdale, has obtained $5\frac{1}{2}$ per cent. more, whilst the proportions indicated by all the other gauges are less than in 1846 and 1847.

It will also be observed that the stations at 1290 and 1334 feet, which in summer receive much less rain than at 1900 feet, in the winter months receive more. This deficiency is obviously owing to the greater proportion of snow deposited at and lost to the instrument at the higher station.

Now, as in the winter months the mountain gauges give no indication of a large proportion of the fall of snow, all of which is secured to the valley stations by their being daily examined, in order to show fairly the gradation from the valley upwards, we must exclude those months, and take in as elements in the calculation, the summer months only.

Annexed are the receipts of the mountain gauges and those of the adjacent valleys, during the summer of 1848:—

	inches.
Stye Head, 1290 feet above the sea	60·35
Seatollar Common, 1334 feet above the sea	57·97
Sparkling Tarn, 1900 feet above the sea	70·95
Great Gabel, 2925 feet above the sea	46·81
Sca Fell Pike, 3166 feet above the sea	49·46
Wastdale, the nearest valley	50·16
Eskdale Head, valley to the S.S.E., $3\frac{1}{2}$ miles distant . . .	37·69
Eskdale, centre of valley to the S.S.E., $5\frac{1}{2}$ miles distant . .	32·46
Ennerdale, valley to the N.W., $3\frac{3}{4}$ miles distant	42·96
Loweswater, valley to the N.N.W., $7\frac{1}{2}$ miles distant . . .	34·52
Buttermere, valley to the N.N.W., $4\frac{1}{2}$ miles distant . . .	44·57
Gatesgarth, valley to the N., $2\frac{1}{2}$ miles distant	57·66

It will be perceived that the increase in the warmer months up to 2000 feet, is great and rapid; and even at the highest attainable elevation in England, the quantity of rain in those months which are free from snow, considerably exceeds the deposit in most of the circumjacent valleys. Indeed (Langdale and Seathwaite excepted) Gatesgarth is the only place which materially exceeds Sca Fell and Gabel in quantity; but as Langdale Head is ten miles distant, and as Seathwaite, besides being several miles to the northward, exceeds enormously the wettest of the other valleys, it is obvious that it would not be fair to institute a comparison between them. If the whole of the snow which falls at the mountain stations could be secured, or an exact equivalent in water be allowed for it, there can be no doubt that the *annual* results would be similar to those for the summer months *only*; but in consequence of the greater proximity of the clouds to the earth in the winter months, the proportions with respect to the valley would probably be somewhat less.

Of late, I have always carried with me a hygrometer of known accuracy on visiting the Lake Districts, and all experiments which I have made on the hygrometrical state of the atmosphere at considerable altitudes above the sea, tend to establish the law which this investigation has brought to light, by showing that the degree of humidity increases upwards from the earth's surface, and that the condition, or combination of conditions most favourable for the condensation and precipitation of vapour in the greatest abundance, does obtain somewhere about 2000 feet above the sea level.

It is probable that the atmosphere is generally, at or near the point of saturation, at and above 2000 feet; but as the air temperature decreases with every further increase of elevation, its capacity for vapour is proportionately diminished, and consequently there will be less to precipitate than at the point where the temperature of the air and that of the dew-point first begin to balance each other.

From the nature of the research, it is quite impossible to obtain regular and connected observations on the hygroscopic state of the atmosphere at such great heights, but in course of time I hope to bring together a sufficient number of data to enable us to connect together some of those links in the great chain of causation which regulates the gradation and amount of precipitation at various altitudes above the earth's surface.

I am fully aware that the physical law indicated by these results, is at variance with the experiments of many careful observers, and with the inferences drawn from them by scientific men of the highest standing. But, with every deference to the opinions and deductions of these eminent authorities, it must be admitted that they have been arrived at from somewhat scanty materials. So far as my knowledge extends, no investigation of any extent or continuity had been made in this department of meteorology previously to that set on foot in the Lake District, about three years ago. The facts previously on record, with few exceptions, referred to comparatively moderate altitudes, mostly under 1000 feet, and as some of the experiments were made on the mountain breasts, the results would vary greatly, according to the gradient and position of the acclivity, and as the gauges were placed either on the windward or leeward side of the hill; regard must also be paid to the season of the year, for a gauge which in summer receives considerably more rain than the valley, may in winter obtain less; and where the instruments are read off at long intervals, there will be no inconsiderable loss from evaporation. Moreover, experience convinces me that little dependence can be placed on the results obtained from gauges stationed on the *side* of a hill, with whatever care they may have been secured. For there cannot be a doubt, that a pluviometer placed on the breast of a mountain, even on the windward side, will receive much less rain than it would do if stationed on a hill-top of equal elevation.

Thus, the gauge at Brant Rigg in Wastdale, about 500 feet above the sea, though on a comparative flat, but with abruptly rising ground behind it, in the summer of

last year has not only received less rain than the valley, but a smaller quantity by 6 inches than at 3166 feet on Sca Fell Pike.

At a future time I hope to follow out this inquiry more fully and systematically, by placing pluviometers at different heights on the breast of Sca Fell, with the view of ascertaining the effect produced by position on rising ground, over or under that due to the respective elevations.

It is not pretended that the law which appears to regulate the distribution of rain in the mountain district of Cumberland, will equally apply to every similar locality; it will doubtless be variously modified according to latitude, position, and many other circumstances; in some situations all trace of the law may disappear, and in others it is possible that it may be reversed.

As my sole object in this inquiry is a search after truth, should my inferences and deductions be found to be incorrect by any one who may investigate the subject more fully and successfully in another locality, I shall feel no hesitation in acknowledging the error.

In addition to the chief objects of research, I record the particulars of all extraordinary phenomena, such as thunder and hail-storms, great floods, and particularly whirlwinds (to which the district is very liable), with a distant prospect of combining the whole in a separate paper, treating of the physical geography of the Lake Country.

Whitehaven, February 6, 1849.