

XXXVI. *Astronomical Observations made at the North Cape, for the Royal Society. By Mr. Bayley.*

- 1769 April 28, 29, and 30, got the observatory and dwelling-house built, and instruments on shore.
- May 1 Set up an oak plank $4\frac{1}{2}$ inches thick, and 14 inches wide. This plank was set a little more than 2 feet in the ground, and well rammed with earth and stones so that it was very steady and firm; to which I screwed the astronomical clock truly perpendicular (by which means it was independant of the observatory and its shaking by the wind) and set it going nearly with sidereal time.
- 3 Set up the transit instrument nearly in the meridian.
- 4 Examined the line of collimation of the quadrant, and found it correct.
Examined the clock, and found the pendulum to vibrate $1^{\circ}\frac{1}{2}$ on each side of nothing.

Here follow some corresponding altitudes of the Sun, from which the going of the clock is determined.

Z. D.

		'	"	h	'	"	'	"			
h	6	27	13	23	34	37	42	22	Morning	Sun's upper limb	} 61,30
		20	58	6	13	30	5	48	Afternoon		
		5	57	0	14	31	23	8	Morning	Sun's upper limb	} 50,0
		42	6	5	33	32	24	55	Afternoon		
O	7	29	16:	0	38	39:	48	24:	Morning	Sun's upper limb	} 57,36 very hazy
		26	24:	5	16	59:	—	—	Afternoon		
		39	20:	0	49	17:	59	44:	Morning	Sun's lower limb	}
		16	18:	—	—	—	—	—	Afternoon		
D	8	29	53	23			44	29	Morning	Sun's upper limb	} 61,20
		34	0	6			19	25	Afternoon		
		37	39	23	45	13	—	—	Morning	Sun's lower limb	}
		26	16	6	18	44	—	—	Afternoon		
h	13	26	46	21	33	0	38	59	Morning	Sun's upper limb	} 71,20
		33	23		39	35	45	34	Afternoon		
<p>This day set up a meridian post nearly in the meridian, by help of the quadrant at about half a mile from the observatory. Examined the pendulum of the clock and found it to vibrate $1^{\circ}\frac{1}{2}$ on each side of nothing.</p>											
D	15	—	—	1	9	25	18	54	Morning	Sun's upper limb	} 55,40
		—	—	5	49	26	39	54	Afternoon		
		9	59	1	19	46	—	—	Morning	Sun's lower limb	}
		48	48	5	39	0	—	—	Afternoon		
h	18	48	56	21	55	4	1	8	Morning	Sun's upper limb	} 70,5
		55	31	22	1	42	—	—	Afternoon	Sun's lower limb	
h	20	—	—	1	36	21	46	20	Morning	Sun's upper limb	} 54,12
		—	—	6	1	48	51	50	Afternoon		
		37	3	1	47	16	—	—	Morning	Sun's lower limb	}
		1	10	5	50	56	—	—	Afternoon		

1769

Corresponding Altitudes of the SUN.

1769

Z. D.

☉	21	-- --	22 6 2	12 6	Morning	Sun's upper limb	} 69,27
		6 28	12 38	-- --	Morning	Sun's lower limb	
♀	26	At 7 ^h 22' per clock, clock stopped, but by what cause I cannot find, it not being down.					
		At 9 ^h I set it going again, as near as I could guess, with sidereal time.					
		Pendulum vibrates 1 ⁰ $\frac{1}{2}$ on each side of nothing.					
24 June	1	Wound up the clock. The pendulum vibrates 1 ⁰ $\frac{1}{2}$ on each side of nothing.					
h	3	49 45	22 55 59	2 3	Morning	Sun's upper limb	} 67,40
		-- --	10 34 52	28 47	Afternoon	Sun's lower limb	
		56 23	23 2 34	-- --	Morning	Sun's upper limb	} 66,50
		34 28	10 28 14	-- --	Afternoon	Sun's lower limb	
		-- --	23 6 16	12 19	Morning	Sun's upper limb	} 63,20
		-- --	10 24 31	18 26	Afternoon	Sun's lower limb	
		6 40	23 12 53	18 51	Morning	Sun's upper limb	} 61,54
		24 4	10 17 52	11 56	Afternoon	Sun's lower limb	
		-- --	23 49 31	55 38	Morning	Sun's upper limb	} 63,20
		-- --	9 41 12	35 5	Afternoon	Sun's lower limb	
		49 56	23 56 14	02 18	Morning	Sun's upper limb	} 61,54
		40 47	9	28 24	Afternoon	Sun's lower limb	
		-- --	0 7 30	13 42	Morning	Sun's upper limb	} 58,42
		-- --	9 23 13	17 2	Afternoon	Sun's lower limb	
		7 55	14 16	20 30	Morning	Sun's upper limb	} 56,40
		22 48	9 16 26	10 12	Afternoon	Sun's lower limb	
		-- --	0 48 51	55 26	Morning	Sun's upper limb	} 55,5
		-- --	8 41 50	35 13	Afternoon	Sun's lower limb	
		49 19	0 56 2	-- --	Morning	Sun's upper limb	} 53,0
		-- --	8 34 37	-- --	Afternoon	Sun's lower limb	
		-- --	1 16 46	23 47	Morning	Sun's upper limb	} 52,5
		-- --	8 13 51	6 50	Afternoon	Sun's lower limb	
		17 14	1 24 26	-- --	Morning	Sun's upper limb	} 50,50
		13 23	8 6 11	-- --	Afternoon	Sun's lower limb	
		-- --	1 39 55	47 29	Morning	Sun's upper limb	} 67,0
		-- --	7 50 41	-- --	Afternoon	Sun's lower limb	
		41 27	1 48 11	-- --	Morning	Sun's upper limb	} 52,5
		-- --	7 42 29	-- --	Afternoon	Sun's lower limb	
		-- --	2 13 44	22 27	Morning	Sun's upper limb	} 50,50
		-- --	7 16 52	8 7	Afternoon	Sun's lower limb	
		14 18	2 23 19	-- --	Morning	Sun's upper limb	} 52,5
		16 18	7 7 18	-- --	Afternoon	Sun's lower limb	
		-- --	2 30 20	39 56	Morning	Sun's upper limb	} 50,50
		-- --	7 0 16	50 39	Afternoon	Sun's lower limb	
		30 56	2 40 52	51 8	Morning	Sun's upper limb	} 67,0
		59 40	6 49 45	39 24	Afternoon	Sun's lower limb	
		-- --	2 56 16	7 58	Morning	Sun's upper limb	} 52,5
		-- --	6 34 20	-- --	Afternoon	Sun's lower limb	
		57 2	3 9 6	-- --	Morning	Sun's upper limb	} 50,50
		33 33	6 21 31	-- --	Afternoon	Sun's lower limb	
24	8	at	11 15 52,5	a mean of four observations Z. D. ☉ L. = 69° 26' 15"			
♀	9	-- --	23 21 46	27 57	Morning	Sun's upper limb	} 67,0
		-- --	10 58 14	52 6	Afternoon	Sun's lower limb	
		22 11	23 28 26	-- --	Morning	Sun's upper limb	} 67,0
		57 49	10 51 39	-- --	Afternoon	Sun's lower limb	

1769

Corresponding Altitudes of the SUN.

Z. D.

	'	"	'	"	'	"			
♀ June 9	33	43	23	0	46	0	Morning	Sun's upper limb	65,32
	46	18	10	40	34	0	Afternoon		
	40	16	23	46	—	—	Morning	Sun's lower limb	65,32
	39	45	10	33	—	—	Afternoon		
	—	—	1	28	—	—	Morning	Sun's upper limb	57,0
	—	—	8	51	—	—	Afternoon		
	29	12	1	36	42	56	Morning	Sun's lower limb	57,0
	50	40	8	43	36	54	Afternoon		
♂ 10	—	—	23	28	34	32	Morning	Sun's upper limb	66,44
	—	—	10	56	—	—	Afternoon		
	28	45	23	34	—	—	Morning	Sun's lower limb	66,44
	0	0	11	0	—	—	Afternoon		
	—	—	0	59	5	50	Morning	Sun's upper limb	59,26
	—	—	9	28	22	22	Afternoon		
	59	56	1	6	—	—	Morning	Sun's lower limb	59,26
	—	—	9	21	—	—	Afternoon		
	—	—	1	20	27	32	Morning	Sun's upper limb	57,48
	—	—	9	7	—	—	Afternoon		
	21	26	1	28	—	—	Morning	Sun's lower limb	57,48
	6	47	9	0	—	—	Afternoon		
♂ 11	—	—	1	42	48	39	Morning	Sun's upper limb	57,30
	—	—	9	27	21	1	Afternoon		
	42	31	1	49	—	—	Morning	Sun's lower limb	57,30
	27	8	9	20	—	—	Afternoon		
	—	—	2	9	16	15	Morning	Sun's upper limb	55,32
	—	—	9	0	—	—	Afternoon		
	9	38	2	16	24	8	Morning	Sun's lower limb	55,32
	0	0	9	0	—	—	Afternoon		
	—	—	3	24	34	14	Morning	Sun's upper limb	50,54 hazy
	—	—	45	8	35	15	Afternoon		
	25	6	3	35	45	48	Morning	Sun's lower limb	50,54 hazy
	44	23	7	34	23	38	Afternoon		
♂ 18	17	16	0	23	29	30	Morning	Sun's upper limb	64,36
	17	13	11	11	4	58	Afternoon		
	23	50	0	30	36	8	Morning	Sun's lower limb	64,36
	10	40	11	4	58	20	Afternoon		
	—	—	0	39	45	17	Morning	Sun's upper limb	63,20
	—	—	0	0	49	16	Afternoon		
	39	37	0	45	—	—	Morning	Sun's lower limb	63,20
	—	—	10	48	—	—	Afternoon		
	—	—	2	22	29	5	Morning	Sun's upper limb	55,25
	—	—	9	12	5	24	Afternoon		
	22	30	2	29	—	—	Morning	Sun's lower limb	55,25
	11	54	9	0	—	—	Afternoon		
	—	—	3	32	41	56	Morning	Sun's upper limb	51,3
	—	—	8	2	52	26	Afternoon		
	32	58	3	42	53	6	Morning	Sun's lower limb	51,3
	1	27	7	51	41	20	Afternoon		

From the above corresponding Altitudes the going of the Clock is determined.

	Apparent noon per clock, per equal al- titudes.	Clock too slow for sidereal time.	Rate of clock with respect to sidereal time.
	h ' "	' "	"
May 6	2 53 31,0	1 1,0	+4,6
8	3 1 25,2	0 52,3	+1,2
15	3 28 59,2	0 43,2	-1,3
20	3 48 43,9	0 49,6	
½ June 3	4 45 5,1	1 13,6	-0,5
9	5 9 46,4	1 16,6	+1,3
10	5 13 56,9	1 15,3	+2,1
11	5 18 7,3	1 13,2	-0,2
15	5 34 42,5	1 14,0	+0,07
18	5 47 10,3	1 14,2	
	Mean rate of clock		+0,91

Clock stopped.

TRANSITS taken with a Transit Instrument over the Meridian, on the Island of Maggerø, or the North Cape of Europe.

1769

$\frac{1}{2}$ May 20 $3^h 49' 30'', 7''$ transit of Sun's center at transit instrument, it being adjusted to the meridian mark, which was put nearly in the meridian. — And $3^h 48' 43'', 9''$ apparent noon per equal altitudes. By this it appears, the meridian mark is west of the true meridian.

\odot 21 Shifted the meridian mark nearer to the true meridian.

	1 Wire.	2 Wire.	3 Wire.	4 Wire.	5 Wire.	
	' "	' "	h ' "	' "	' "	
$\frac{1}{2}$ June 3		43 $11\frac{1}{2}$	4 44 $1\frac{1}{2}$			\odot 1 L.
			46 17	47 $7\frac{1}{2}$		\odot 2 L.
$\frac{1}{2}$ 9			5 10 58,6			\odot 2 L. passed the merid.
	11 $\frac{5}{8}$	27 8+	18 28 5—	29 2—	58—	α Lyræ.
	59—	43 $\frac{3}{8}$	19 38 28+	13 $\frac{3}{8}$	58	α Aquilæ.
$\frac{1}{2}$ 10	47 8	47 54	10 48 39	49 $24\frac{1}{2}$		δ 1 L.
			12 45 14			Polaris S. P.
	32 v	19—	14 4 6—	53 $\frac{1}{2}$		Arcturus.
	14 v	11—	18 28 7+	4+	0 $\frac{5}{8}$	α Lyræ.
\odot 11	15 24—	16 12+	5 17 0 $\frac{1}{2}$			\odot 1 L.
			19 18 $\frac{1}{2}$	20 $7\frac{5}{8}$	20 55 $\frac{1}{2}$	\odot 2 L.
$\frac{1}{2}$ 15		32 46 $\frac{1}{2}$	5 33 34 $\frac{1}{2}$			\odot 1 L.
			35 53+	36 41 $\frac{1}{2}$		\odot 2 L.
$\frac{1}{2}$ 16	17 52+	18 39—	4 19 26—	20 13+		$\frac{1}{2}$ 2 L.
$\frac{1}{2}$ 17	40 18	41 6+	5 41 54+			\odot 1 L.
			44 15 $\frac{1}{2}$	45 2 $\frac{1}{2}$	45 50 $\frac{5}{8}$	\odot 2 L.
	33+	3 20 $\frac{1}{2}$	14 4 7 $\frac{1}{2}$	4 55		Arcturus.
	42 22+	43 11—	17 43 59+		47 54+	\odot 1 L.
			46 17 $\frac{1}{2}$	47 5 $\frac{5}{8}$		\odot 2 L.
	26 14	27 11	18 28 7 $\frac{1}{2}$	4 $\frac{1}{2}$		α Lyræ.
		17 31—	4 10 17 $\frac{1}{2}$	19 4 $\frac{3}{8}$		$\frac{1}{2}$ 2 L.
\odot 18	44 28—		5 46 24 $\frac{1}{2}$			\odot 1 L. } flying clouds.
		47 35—	48 23 $\frac{1}{2}$	49 12 $\frac{1}{2}$		\odot 2 L. }
			12 45 30:			Polaris S. P.

☉ 1 L.
 ☉ 2 L.
 ☉ 2 L. passed the merid.
 α Lyræ.
 α Aquilæ.
 δ 1 L.
 Polaris S. P.
 Arcturus.
 α Lyræ.
 ☉ 1 L.
 ☉ 2 L.
 ☉ 1 L.
 ☉ 2 L.
 ♀ 2 L.
 ☉ 1 L.
 ☉ 2 L.
 Arcturus.
 ☉ 1 L.
 ☉ 2 L.
 α Lyræ.
 ♀ 2 L.
 ☉ 1 L.
 ☉ 2 L. } flying clouds.
 Polaris S. P.

TRANSITS compared with equal Altitudes, for finding the Error of the Meridian Mark.

		Transits of Sun at transit instrument.	Noon per equal altitudes.	Differences.
		h ' "	h ' "	"
June	3	4 45 9.3	4 45 5.1	4.2 W.
	11	5 18 9.9	5 18 7.3	2.6 W.
	15	5 34 43.9	5 34 42.5	1.4 W.
	18	5 47 13.0	5 47 10.3	2.7 W.
		Mean difference		2.7 West.

Hence it appears, that the Sun passed at the transit instrument too late 2".7 by a mean of these observations, which shews the mark was west of the true meridian; which gives the azimuth of mark = 50" south westerly.

TRANSIT of VENUS, observed at the NORTH CAPE.

¹⁷⁶⁹
June 3 At 13^h 46' 40" per clock, or 9^h 0' 2", apparent time, the Sun came out from under a cloud, with Venus on it, about $\frac{1}{4}$ th of her diameter; and at 14^h 0' 41" or 9^h 14' 1", apparent time, Venus's outer limb seemed to be in contact with the Sun's limb; but no light of that part of the Sun's limb could be seen, Venus being apparently joined to the Sun's limb by a black ligament, which gradually diminished in breadth; and at 14^h 1' 36", or 9^h 14' 56", the Sun's light broke through it, and Venus and the Sun were to appearance perfect (this was certain to about 10 or 15 seconds of time), and the black ligament contracted itself, so that Venus was considerably within the Sun's limb, suppose $\frac{1}{2}$ th of her diameter. During these observations the air was red and hazy, and the Sun's limb very tremulous, and the spots in the Sun very indistinct, and Venus seemed very ill defined when on the Sun. But a better idea will be formed of the bad appearance of Venus at the internal contact, owing to the very hazy state of the air, from the representation of it, plate XIII.

Here follow some measures I was able to take during the time the clouds were off the Sun.

Time per clock.	App. time.	Measures of microm. in inches, &c.	Measures of microm. in degrees, &c.
--------------------	------------	--	---


Soon after the internal contact, measures of Venus's diameter.

	0,1 24
	0,1 23 +
	0,1 23 —
	0,1 23 +
	0,1 22
	0,1 22
	0,1 23 —
	0,1 24 —
	0,1 23 —
	0,1 23
	0,1 23 —
	0,1 23 —
Mean of	12
	0,1 22,8 = 55",32

Measures of the Sun's horizontal and vertical diameters, taken immediately after Venus's diameter.

Horizontal Diameter.		Vertical Diameter.	
4,9 $\frac{1}{2}$	21 $\frac{1}{2}$	4,8 $\frac{1}{2}$	12
4,9 $\frac{1}{2}$	23	4,8 $\frac{1}{2}$	10
4,9 $\frac{1}{2}$	23	4,8 $\frac{1}{2}$	11
4,9 $\frac{1}{2}$	21 $\frac{1}{2}$	4,8 $\frac{1}{2}$	9
Mean	4,9 $\frac{1}{2}$ 22,2	31' 37",61	4,8 $\frac{1}{2}$ 10,5 30' 50",73

Difference



**Difference of Declinations of the North Limbs
of the Sun and Venus.**

Time per clock.			App. time.			Measures of microm. in inches, &c.	Measures of microm. in degrees, &c.
h	'	"	h	'	"		' "
14	38	37	9	51	50	$0,5\frac{1}{2}$ 1	3 29,7
	44	30		57	32	$0,5\frac{1}{2}$ 9	3 35,8
	46	56	10	0	8	$0,5\frac{1}{2}$ 13	3 38,8
	52	45		5	56	$0,5\frac{1}{2}$ 23	3 46,4

**Difference of Declination of the South Limbs
of the Sun and Venus.**

14	54	50	10	8	1	$4,2\frac{1}{2}$ 5	26 58,6
	57	43		10	53	$4,2\frac{1}{2}$ 0	26 54,8
	59	52		13	2	$4,2$ 20	26 51,0
15	2	0		15	9	$4,2$ 16	26 48,0

**Equatoreal Distances of the Western Limbs of
the Sun and Venus.**

15	8	5	10	21	13	3,1 17	19 50,8
	10	28		23	36	3,1 2	19 39,4
	12	15		25	23	$3,0\frac{1}{2}$ 20	19 34,1
	13	48		26	56	$3,0\frac{1}{2}$ 12	19 27,9

**Equatoreal Distances of the Eastern Limbs of
Venus and the Sun.**

15	15	50	10	28	57	$6,8\frac{1}{2}$ $8\frac{1}{2}$	11 49,4
	17	38		30	45	$1,8\frac{1}{2}$ 20	11 58,1
	19	33		32	40	1,9 8	12 8,0
	22	0		35	6	1,9 19	12 16,4
	24	5		37	11	$1,9\frac{1}{2}$ 7	12 26,2

**Greatest and least Distances of Venus's nearest
Limbs from the Sun's Limbs, for finding
the nearest Distance of their Centers.**

15	49	3	11	2	4	$0,6\frac{1}{2}$ 17	4 19,9
	51	22		4	23	4,0 2+	25 21,3
	54	24		7	25	0,7 7	4 31,3
	58	1		11	1	$3,9\frac{1}{2}$ 10	25 8,4
16	0	40		13	39	$3,9\frac{1}{2}$ 7	25 6,1
	4	4		17	3	0,7 23	4 43,4
	6	24		19	22	$0,7\frac{1}{2}$ 5	4 48,8
	9	55		23	53	3,9 14	24 52,5
	16	29		29	26	0,7 12	4 35,1
	21	31		34	27	0,7 8	4 32,1
16	25	55	11	38	51	$3,8\frac{1}{2}$ 20::24	38,0::

**A Table for reducing the Scale of
the micrometer to Degrees, &c.**

Inches.	Value.	Tenths.	Value.	V.	Value.
'	"	'	"		"
1	6 19,95	1	0 37,99	1	0,76
2	12 39,91	2	1 15,99	2	1,52
3	18 59,86	3	1 53,98	3	2,28
4	25 19,81	4	2 31,98	4	3,04
5	31 39,77	5	3 9,97	5	3,80
		6	3 47,97	6	4,56
		7	4 25,97	7	5,32
		8	5 3,96	8	6,08
		9	5 41,96	9	6,84
				10	7,60
				11	8,36
				12	9,12
				13	9,88
				14	10,64
				15	11,40
				16	12,16
				17	12,92
				18	13,68
				19	14,44
				20	15,20
				21	15,96
				22	16,72
				23	17,48
				24	18,24
				25	18,99

On examination, the micro-
meter scale wanted no ad-
justment.

ECLIPSE of the SUN, observed at the NORTH CAPE.

1769

June 3 At 1^h 48' 4", the clouds clearing away, I saw the Sun, and the Moon had made a small impression or notch in the Sun's limb; by observing the increase of the eclipse, I suppose it began 4, 5, or 6 seconds sooner than I first saw it, or at 1^h 48' 0" per clock, or 20^h 59' 19" apparent time, nearly.

Distances of the Cusps.

Time p clock	App. time	Measures	Reduced
h / "	h / "		
2 10 22	21 21 0	3,7 5	23 29,6
13 19	24 34	3,8½ 9	24 29,6
19 1	30 15	4,1½ 6	26 21,3
21 3	32 16	4,2 7	26 41,1
23 50	35 3	4,3½ 8	27 38,8
27 38	38 50	4,4½ 14	28 21,4
31 10	41 21	4,5½ 10	28 56,4
34 56	46 7	4,6 13	29 17,7
36 4	47 14	4,6½ 15	29 38,2
39 11	50 21	4,7 0	29 45,8

Sun's horizontal Diameter, measured directly after the Eclipse ended.

4,9½ 21½
4,9½ 20
4,9½ 20
4,9½ 22+
4,9½ 21—

Mean 4,9½ 20,9=31' 35",9

Measures of the lucid Part, near the Middle of the Eclipse.

2 43 7	21 54 16	1,1 5	7 1,7
44 11	55 20	0,9½ 22	6 17,7
47 50	58 59	0,9 24	6 0,2
49 35	22 0 43	0,9 9	5 48,4
52 14	3 22	1,0½ 11	6 47,3

Distances of the Cusps.

2 54 55	22 6 2	4,6½ 19	29 41,2
57 1	8 8	4,7 3	29 48,1
59 55	11 2	4,6 14	29 18,4
3 2 24	13 30	4,6 10	29 15,4
4 31	15 37	4,6 7	29 13,1
6 37	17 42	4,5½ 9	28 55,6
8 13	19 18	4,5 20	28 45,0
10 34	21 38	4,4½ 18	28 24,5
14 57	22 26 1	4,3 17	27 26,7

By comparing the end with Mr. Maskelyne's observation at Greenwich, the difference of meridians comes out 1^h 44' 6" of time. = 26° 1' 30" east, or difference of longitude of my observatory from Greenwich to the east. From whence the longitude of the point called the North Cape is 25° 49' east of Greenwich.

Clouds came on, so that I saw the Sun no more till 3^h 38' 0" per clock, and it broke away very clear, and continued clear to the end, which was at 3^h 48' 19" per clock, or 22^h 59' 17" apparent time. The air being very clear, the end seemed certain to about two seconds.

The telescope used was a reflector of 2 feet focus, made by Mr. Dollond; and the magnifying power, applied for the ingress of Venus, and the beginning and end of the solar eclipse, was 100. The magnifying power used with the micrometer, was 50.

Adjustments of the telescope with the different eye-pieces and little speculums, as combined together when used. N. B. — shews that (o) on the vernier, is behind, or to the left hand of the first division on the scale; and + to the right hand, or before.

Long eye-piece and short focus little speculum magnifying 100 times = $-0 + 11\frac{1}{2}$ on vernier.

Long eye-piece and long focus little speculum with micrometer on $= -0 + 19\frac{2}{3}$ on vernier

The eye-piece with moveable wires, and long focus little speculum, and micrometer } = + 0 + 4 on vernier.

These are a mean of 10 or 12 observations each.

The Value of the Scale of the Object Glass Micrometer was found as follows :

A bafe of 120 feet was carefully measured on level ice (which was covered with frozen snow about half an inch thick), with two 10 feet fir rods; and the measure being taken four different times, no one differed from any of the others so much as $\frac{1}{100}$ th of an inch. But as the fir rods might not consist of 10 feet exactly of the same standard from which the micrometer scale was laid off, I therefore took six inches between the fine points of a pair of compasses, from the micrometer scale; and, by repeating this measure, found the exact length of the fir rods in measures of the micrometer scale; and thence corrected the length of the bafe, found immediately by the rods, and reduced to the standard of the micrometer scale. Perpendicularly over the bafe, at one end, was placed the center of the divided object glass, and perpendicularly over the bafe, at the other end, a board was placed, having its plane at right angles with the bafe line, white paper pasted on its upper part; and at the height of the center of the object glass, nearly above the horizon, was a strong black line drawn, as $AB = 8$ inches, Ag and Ae being small but equal distances from A ; by bringing (ed and gb) to coincide alternately with (Aa) the error of adjustment of the micrometer scale was found; and by separating the glasses till (Aa) and (Bb) made but one line; from this separation the value of one extent of the micrometer scale was found, by making this proportion. As the length of the bafe is to the length (AB) :: so is 206265'' (the number of seconds in an arch equal to the radius of a circle) to a fourth number, which will be the seconds of the angle measured by that opening of the glasses, which is shewn by the micrometer scale; and other angles will be in direct proportion to the respective measures when the table was formed.

ZENITH DISTANCES taken with an Astronomical Quadrant, of one Foot Radius, made by Mr. Bird, at the NORTH CAPE of EUROPE.

		Bar. Ther.					
		Inches.	out	in			
			o	o			
1769							
☉ May	14	29,80	43	40	☉ U. L.	on the merid.	
	20	29,71	45	41	☉ L. L.	ditto	
♂ June	3	29,93	56	58	☉ L. L.		
					☉ U. L.		
	11	29,57	46	42	☉ L. L.		
					☉ U. L.		
	15	29,66	41	40	☉ U. L.		
					☉ L. L.		
	16	29,70	44	39	♀ Center		
	17	29,72	44	42	☉ L. L.		
					☉ U. L.		
		29,71	47	48 $\frac{1}{2}$	Arcturus		
		29,70	52	51 $\frac{1}{2}$	☉ L. L. } on the north		
					☉ U. L. } meridian.		
		29,70	52	51 $\frac{1}{2}$	α Lyrae		
		29,18	52	54 $\frac{1}{2}$	♀ Center		

Interior arch				Exterior arch				Exterior arch reduced			
o	′	″		o	′	″	″	o	′	″	
51	59	10		55	1	26	15	51	58	59,2	
51	10	40,0		54	2	12	8	51	10	45,9	
48	52	44,0		52	0	18	16	48	52	36,6	
48	20	54,0		51	2	9	0	48	20	49,8	
				51	1	11	12	48	7	26,7	
47	35	37,0		50	3	3	0	47	35	40,3	
				50	2	5	8	47	22	41,3	
47	54	27,8		51	0	13	14	47	54	33,8	
52	13	36+		55	2	27	17	52	13	27,4	
47	50	28,0		51	0	4	0	47	50	30,5	
47	18	50,0		50	1	28	10	47	18	42,0	
50	36	8,0		53	3	30	10	50	36	21,2	
				91	1	12	24	85	37	41,1	
85	7	29+		90	3	6	0	85	7	19,4	
32	25	18+		34	2	11	0	32	25	22,0	
52	27	0,0		55	3	25	0	52	26	55,4	
								From			

From the foregoing Zenith Distances of the Sun, and of Arcturus, and α Lyræ, the Latitude of the Observatory is determined, as follows :

		From	Latitude		
			°	'	"
1769					
May	14	⊙ U. L.	71	0	43,2
	20	⊙ L. L.	71	1	0,1
June	3	⊙ Center	71	0	43,5
	11	⊙ Center	71	0	39,8
	15	⊙ Center	71	0	39,7
	17	⊙ Center	71	0	40,6
		Arcturus	71	1	0,9
		α Lyræ	71	0	48,6
Mean - - -			71	0	47,0

From whence the latitude of the point of land called the North Cape is $71^{\circ} 10'$ north.

By a great many trials with a very good compass, of Dr. Knight's construction, I found the variation to be 6 degrees west ; and by a dipping needle, I found, by repeated trials, the dip of the north end of the needle to be 79 degrees.

May 15, at $13^h \frac{1}{4}$ P. M. apparent time, or $1^h 7'$ after high water, by a mean of 7 observations, I found the dip of the horizon of the sea, from the observatory, to be $12' 18''$. Height of the barometer 29,70 inches ; thermometer, without, 24° ; thermometer, within, 28° . And May 20, at $7^h \frac{3}{4}$ P. M. apparent time, or $7^h 33'$ after high water, from a mean of 8 observations, I found the dip = $12' 25''$, 5 ; barometer 29,70 inches ; thermometer, without, 43° , and, within, 40° . Both these observations were made on the N. N. E. point of the true compass. During each of these observations the water was very smooth, and the horizon clear.—I found it was high water, at the full and change of the Moon, at $3^h 44'$ P. M. apparent time, at the Cape ; and, by a series of observations, I found the water to rise 8 feet 1 inch, nearly, perpendicular at the spring tides ; and at neap tides 6 feet 8 inches, perpendicular ; and the tides seemed to follow very regular, as they ought to do when not disturbed by bad weather.—June 8, I found the height of the observatory 140 feet 6 inches above low water mark.

A JOURNAL of the Barometer and two Thermometers, during the Time I was on Shore in the Island of MAGGERØ, or NORTH CAPE, viz. from the 1st of May to the 21st of June.

At noon.				
		Bar.	Ther.	
			out	in
		Inches	°	°
1769				
May	1	29,72	19	24
	2	29,68	26	28
	3	29,90	18	22
	4	29,96	28	$31\frac{1}{2}$
	5	29,92	34	32
	6	29,91	58	38
	7	29,74	60	40
	8	29,84	56	39

At noon.

At midnight.

Bar. Ther.
out in

Bar. Ther.
out in doors

Inches

Inches

1769						
May	9	29,90	47	36		
	10	29,97	46	39	29,98	34 36
	11	29,97	39	34	29,96	30 32
	12	29,97	30	31	29,97	27 29
	13	30,03	42	36	30,04	34 37
	14	29,80	43	40	29,74	33 37½
	15	29,73	41	37	29,70	28 30
	16	29,66	29	31	29,64	26 28
	17	29,62	51	42	29,60	39 43
	18	29,48	53	45	29,39	38 41
	19	29,34	51	47	29,57	40 40
	20	29,71	45	41½	29,77	37 41
	21	29,51	54½	47	29,46	37 39
	22	29,51	41	38	29,64	36 43
	23	29,77	40	37	29,78	36 36½
	24	29,65	44	41½	29,47	40 40
	25	29,42	52	45	29,71	44½ 45
	26	29,77	44	45	29,78	40 42
	27	29,76	48	44	29,73	36 38
	28	29,70	44	36½	29,66	34 35
	29	29,47	38	33	29,51	38 40½
	30	29,61	50	43	29,61	52 50
	31	29,61	52	48	29,70	42 37
June	1	29,67	41	38	29,72	38 40
	2	29,92	54	50	29,90	48 52
	3	29,93	66	58	29,78	36 36½
	4	29,79	43	43	29,70	46 40
	5	29,77	61	55	29,69	47 49
	6	29,51	47	43	29,50	43 43
	7	29,83	43	47	29,90	42 44
	8	29,89	47	45	29,90	41 40
	9	29,84	49	47½	29,74	40 42
	10	29,67	48	42	29,66	41 43
	11	29,57	46	42	29,58	33 38
	12	29,48	38	35	29,51	31½ 33
	13	29,56	38	34	29,52	35 34
	14	29,53	39	36	29,57	35 35½
	15	29,66	41	40	29,64	34 35
	16	29,51	38	38	29,62	38 37½
	17	29,72	44	42	29,70	52 51½
	18	29,18	52	54½	29,52	44 46
	19	29,42	37	39	29,38	36 38
	20	29,67	36	37	29,49	32 34
	21	29,54	36	38	29,60	31½ 33
	22	29,46	35	37		

The thermometers were kept, the one in the observatory, and the other without, in the open air; but always in the shade; and were always observed at noon and midnight, after the 10th of May; but before, only at noon.

The INSTRUMENTS used at the NORTH CAPE, by Mr. BAYLY,
were as follows:

A quadrant of 1 foot radius, and two thermometers, made by Mr. Bird.

A 2 feet reflector, with an achromatic object glass micrometer, by Mr. Dollond.

A transit instrument, of 4 feet, made by Mr. Bird; with an achromatic object glass,
by Mr. Dollond.

A barometer, by Mr. Ramsden.


An astronomical clock with a gridiron pendulum, a journeyman clock, and an alarm
clock, by Mr. Shelton.

A dipping needle, belonging to the Royal Observatory, made by Mr. Graham.

A like set, exclusive of a dipping needle, was used by Mr. Dixon, at the island of
Hammerfest.

N. B. The adjoining chart, and views of the sea-coast and islands, near the North Cape of
Europe, Tab. XIV. were drawn from the joint observations of Messieurs Dixon and
Bayly.

71



T H E N O

THE NORTH CAPE

The Mother & Daughters

Jelmsby I.

Haws I.

THE ISLE OF
M A G G E R O E

The Observatory

Kelwic

Hannings Bay

D A N I S H
F U E R

D
R
T
H

S
E

A


atory
Kelwick

L
A
P
L
A
N
D

The North Cape, S.E. 2 Miles

72.

71



O R

S I

I S

North

By JE




A
C H A R T
of the
S E A C O A S T
and
S L A N D S
near the
North Cape of Europe

By JERE. DIXON and W^m BAYLY 1769.

25


26'



The Isle of Maggeroe .



Jelmsby .



Ingan .

Shipholm .

